

On the Symmetry of the Diffusion Coefficient in Asymmetric Simple Exclusion.

Michail Loulakis¹

ABSTRACT: We prove the symmetry of the diffusion coefficient that appears in the fluctuation-dissipation theorem for asymmetric simple exclusion processes.

Mathematics Subject Classification (2000): Primary 60K35; secondary 82C22

Keywords and Phrases: Diffusion Coefficient, Fluctuation-Dissipation Theorem, Simple Exclusion.

1 Introduction

The fluctuation-dissipation theorem has been an instrumental tool in the analysis of non-gradient interacting particle systems. In the context of asymmetric simple exclusion processes (ASEP) it has been used to establish the hydrodynamic limit for the mean-zero ASEP⁷, as well as the diffusive incompressible limit,² the first order corrections to the hydrodynamic limit,³ the equilibrium fluctuations of the density field,¹ and the diffusive hydrodynamic limit when the initial density profile is constant along the direction of the drift,⁶ for the general ASEP in $d \geq 3$.

The fluctuation-dissipation theorem consists of decomposing the (normalized) particle currents $(w_i)_{1 \leq i \leq d}$ (which are not of gradient form) into gradients of the occupation variable and a rapidly fluctuating term. With a suitable interpretation it can be formulated in the following equation:

$$w_i = \sum_{j=1}^d D_{ij}(\eta(0) - \eta(e_j)) + Lu_i. \quad (1)$$

The matrix $D = (D_{ij})_{1 \leq i, j \leq d}$ is called the *diffusion coefficient* and it naturally appears in the PDEs that arise in the hydrodynamic limit. Explicit and variational formulae for D are available⁴ and it is known to be a smooth function of the particle density.⁵ In this article we prove that D is symmetric, thus answering the question raised by Landim, Olla and Yau in ref. 3 and 4.

2 Notation and Results

Let us fix a finite range probability measure $p(\cdot)$ on $\mathbb{Z}^d$, with $p(0) = 0$. We denote by L the generator of the simple exclusion process associated to $p(\cdot)$. L acts on local functions on the state space $\mathbb{X} = \{0, 1\}^{\mathbb{Z}^d}$ according to:

$$Lf(\xi) = \sum_{x, y} p(y - x) \xi(x) (1 - \xi(y)) (f(\xi^{x, y}) - f(\xi)) \quad (2)$$

¹Statistical Laboratory, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, United Kingdom. E-mail: michail@statslab.cam.ac.uk

where:

$$\xi^{x,y}(z) = \begin{cases} \xi(z) & \text{if } z \neq x, y, \\ \xi(x) & \text{if } z = y, \\ \xi(y) & \text{if } z = x. \end{cases}$$

The symmetric and the anti-symmetric part of $p(\cdot)$ will be denoted by $a(\cdot)$ and $b(\cdot)$ respectively:

$$a(x) = \frac{p(x) + p(-x)}{2}, \quad b(x) = \frac{p(x) - p(-x)}{2}.$$

In order to avoid degeneracies we will assume that the random walk in $\mathbb{Z}^d$ with one step transition probabilities $a(y-x)$ is irreducible, i.e. $\{x : a(x) > 0\}$ generates the group $\mathbb{Z}^d$. An equivalent formulation of this assumption is that the matrix $S = (S_{ij})_{1 \leq i, j \leq d}$ defined by $S_{ij} = \frac{1}{2} \sum p(z) z_i z_j$ is invertible.

The symmetric part of the generator (denoted by L^s) is given by (2) with $p(\cdot)$ replaced by $a(\cdot)$. The measures μ_ρ ($0 \leq \rho \leq 1$), defined as Bernoulli products of parameter ρ over the sites of $\mathbb{Z}^d$ are invariant under the dynamics. We will denote expectations under μ_ρ by $\langle \cdot \rangle_\rho$ and inner products in $L^2(\mu_\rho)$ by $\langle \cdot, \cdot \rangle_\rho$.

The adjoint of L in $L^2(\mu_\rho)$ is the generator L^* of the simple exclusion process associated to the law $p^*(x) = p(-x)$. Local functions form a core of both L and L^* , and thus L^s extends to a self-adjoint operator in $L^2(\mu_\rho)$.

The particle current along the direction e_i is given by:

$$W_i = \frac{1}{2} \sum_z p(z) z_i \xi(0) (1 - \xi(z)) - p(-z) z_i \xi(z) (1 - \xi(0)). \quad (3)$$

Equation (1) is to be understood in the Hilbert space of fluctuations, which we define next. Let $\mathcal{G}_\rho$ be the space of local functions g such that:

$$\langle g \rangle_\rho = 0 \quad , \text{ and } \quad \left. \frac{d}{d\theta} \langle g \rangle_\theta \right|_{\theta=\rho} = 0.$$

For a $g \in \mathcal{G}_\rho$ we define $\tau_x g = g(\tau_x \xi)$, where $\tau_x \xi(z) = \xi(x+z)$. For any $f \in \mathcal{G}_\rho$ and $i \in \{1, \dots, d\}$ we define:

$$\langle g, f \rangle_{\rho,0} := \sum_x \langle g, \tau_x f \rangle_\rho, \quad t_i(g) = \langle g, \sum_x x_i \xi(x) \rangle_\rho.$$

Set $\chi(\rho) = \rho(1-\rho)$ and define

$$\langle \langle g \rangle \rangle_\rho = \sup_{\alpha \in \mathbb{R}^d} \left(2 \sum_{i=1}^d \alpha_i t_i(g) - \chi(\rho) \alpha \cdot S \alpha \right) + \sup_{f \in \mathcal{G}_\rho} (2 \langle g, f \rangle_{\rho,0} - \langle f, (-L^s) f \rangle_{\rho,0}), \quad (4)$$

The Hilbert space of fluctuations $\mathcal{H}(\rho)$ is defined as the closure of $\mathcal{G}_\rho$ under $\langle \langle \cdot \rangle \rangle_\rho^{1/2}$. If we denote by $\mathcal{H}_0$ the space generated by gradients of the occupation variable: $\mathcal{H}_0 = \{\sum \alpha_i (\xi(e_i) - \xi(0)); \alpha \in \mathbb{R}^d\}$, then by Theorem 5.9 in ref. 2 we have:

$$\mathcal{H}(\rho) = \overline{\mathcal{H}_0 + L \mathcal{G}_\rho}. \quad (5)$$

Notice that unless $\sum z p(z) = 0$ the currents W_i do not belong to the space $\mathcal{G}_\rho$. Therefore we define the *normalized* currents $w_i \in \mathcal{G}_\rho$ by:

$$w_i = W_i - \langle W_i \rangle_\rho - (\xi(0) - \rho) \left. \frac{d}{d\theta} \langle W_i \rangle_\theta \right|_{\theta=\rho}. \quad (6)$$

According to (5) there exist coefficients $(D_{ij})_{1 \leq i, j \leq d}$ (which depend on ρ) such that:

$$w_i - \sum_{j=1}^d D_{ij} \times (\xi(0) - \xi(e_j)) \in \overline{L\mathcal{G}_\rho}. \quad (7)$$

The matrix $D = (D_{ij})_{1 \leq i, j \leq d}$ is called the diffusion coefficient of the simple exclusion process. Landim, Olla and Yau proved explicit and variational formulae for D in ref. 4. In the same paper, as well as in ref. 3, the authors question whether there exists a choice of $p(\cdot)$ such that D is asymmetric. The result of this article is the following theorem:

Theorem 1 The diffusion coefficient D defined in (7) is always a symmetric matrix.

3 Some Properties of $\mathcal{H}(\rho)$

In this section we review some properties of the Hilbert space of fluctuations that will be useful in the proof of Theorem 1. We begin with the following lemma.

Lemma 1 If $g \in \mathcal{G}_\rho$ and $h \in \mathbb{Z}^d$, then $\tau_h g = g$ in $\mathcal{H}(\rho)$.

Proof: In view of (4) it suffices to show that:

$$(i) \quad \langle \tau_h g - g, f \rangle_{\rho, 0} = 0, \forall f \in \mathcal{G}_\rho, \quad (ii) \quad t_i(\tau_h g - g) = 0, i = 1, \dots, d.$$

Using the translation invariance of μ_ρ property (i) follows immediately, while

$$\sum_x \langle \tau_h g - g, x_i \xi(x) \rangle_\rho = h_i \sum_x \langle g, \xi(x) \rangle_\rho.$$

The last expression is trivially zero if $\rho \in \{0, 1\}$, while otherwise by differentiating with respect to θ both sides of the following identity

$$\langle g \rangle_\theta = \int g(\xi) \prod_{x \in \text{supp}(g)} \left(\frac{\theta}{\rho} \right)^{\xi(x)} \left(\frac{1-\theta}{1-\rho} \right)^{1-\xi(x)} d\mu_\rho(\xi),$$

we get

$$\sum_x \langle g, \xi(x) \rangle_\rho = \rho \langle g \rangle_\rho + \chi(\rho) \left. \frac{d}{d\theta} \langle g \rangle_\theta \right|_{\theta=\rho} = 0, \quad (8)$$

thus establishing (ii). $\square$

The following lemma is a generalisation of (5.1) in ref. 4 to the general ASEP, and can be proved by polarization of (4). The details are left to the reader.

Lemma 2 Let $g, f \in \mathcal{G}_\rho$ and set $\nabla_{e_k} \xi(0) = \xi(0) - \xi(e_k)$ for $k = 1, \dots, d$. Then:

$$\begin{aligned} (i) \quad & \langle \langle \nabla_{e_k} \xi(0), \nabla_{e_\ell} \xi(0) \rangle \rangle_\rho = \chi(\rho) (S^{-1})_{k\ell}, \\ (ii) \quad & \langle \langle \nabla_{e_k} \xi(0), Lg \rangle \rangle_\rho = -\langle \langle \nabla_{e_k} \xi(0), L^* g \rangle \rangle_\rho = \sum_{\ell=1}^d (S^{-1})_{k\ell} \langle w_\ell, g \rangle_{\rho, 0}, \\ (iii) \quad & \langle \langle \nabla_{e_k} \xi(0), L^s g \rangle \rangle_\rho = 0, \\ (iv) \quad & \langle \langle L^s g, f \rangle \rangle_\rho = -\langle g, f \rangle_{\rho, 0}. \end{aligned}$$

4 The Diffusion Matrix

Recall the definition of the normalised currents w_i given in (3) and (6). It follows by elementary algebra and Lemma 1 that $w_i = W_i^s - h_i$, where

$$W_i^s(\xi) = \frac{1}{2} \sum z_i a(z) (\xi(0) - \xi(z))$$

is the current of the symmetric simple exclusion with generator L^s , and

$$h_i(\xi) = \sum_z z_i b(z) (\xi(0) - \rho) (\xi(z) - \rho).$$

Hence, the normalized currents for the reversed process are given by $w_i^* = W_i^s + h_i$. Let now $C(\rho)$ (resp. $C_*(\rho)$) be the real vector space generated by the currents $\{w_i; i = 1, \dots, d\}$ (resp. $\{w_i^*; i = 1, \dots, d\}$).

We define the linear operator T (resp. T^*) on $C(\rho) + L\mathcal{G}_\rho$ (resp. $C_*(\rho) + L^*\mathcal{G}_\rho$) by:

$$\begin{aligned} T\left(\sum_{i=1}^d \alpha_i w_i + Lg\right) &= \sum_{i,k=1}^d \alpha_i S_{ik} \nabla_{e_k} \xi(0) + L^s g, \\ T^*\left(\sum_{i=1}^d \alpha_i w_i^* + L^*g\right) &= \sum_{i,k=1}^d \alpha_i S_{ik} \nabla_{e_k} \xi(0) + L^s g. \end{aligned}$$

By Theorem 5.9 in ref. 2 we have: $\mathcal{H}(\rho) = \overline{C(\rho) + L\mathcal{G}_\rho} = \overline{C_*(\rho) + L^*\mathcal{G}_\rho}$. Now, just as in Lemma 5.4 in ref. 4, T and T^* are norm bounded by 1, hence they can be extended to $\mathcal{H}(\rho)$. Furthermore, it follows easily by computations based on Lemma 2 that T^* is the adjoint of T with respect to $\langle\langle \cdot, \cdot \rangle\rangle_\rho$ and $T^* \nabla_{e_k} \xi(0)$ is orthogonal to $\overline{L\mathcal{G}_\rho}$. Hence by (7):

$$\langle\langle w_i, T^* \nabla_{e_k} \xi(0) \rangle\rangle_\rho = \sum_{i=1}^d D_{ij} \langle\langle \nabla_{e_j} \xi(0), T^* \nabla_{e_k} \xi(0) \rangle\rangle_\rho,$$

and thus by Lemma 2(i):

$$\chi(\rho) I_d = D \cdot Q$$

where the matrix $Q = (Q_{jk})_{1 \leq j, k \leq d}$ is given by:

$$Q_{jk} = \langle\langle T \nabla_{e_j} \xi(0), \nabla_{e_k} \xi(0) \rangle\rangle_\rho.$$

We are now ready to proceed with the proof of Theorem 1.

Proof (of Theorem 1): Let us denote the reflection operator on $\mathbb{X}$ by

$$R\xi(z) = \xi(-z).$$

The action of R is naturally extended to functions as $Rf(\xi) = f(R\xi)$. Clearly, $R^2 = 1$. Furthermore, the following commutation relation can be readily verified:

$$RL = L^*R. \tag{9}$$

In particular R commutes with L^s and hence, R preserves inner products in $\mathcal{H}(\rho)$.

Notice that W_i^s are anti-symmetric under R , while h_i are R -symmetric. Thus,

$$Rw_i(\xi) = -W_i^s(\xi) - h_i(\xi) = -w_i^*(\xi). \tag{10}$$

It is a direct consequence of (9), (10), and the observation that $R\nabla_{e_k}\xi(0) = -\nabla_{e_k}\xi(0)$ in $\mathcal{H}(\rho)$ that

$$RT = T^*R.$$

Therefore,

$$\begin{aligned} Q_{jk} &= \langle\langle RT\nabla_{e_j}\xi(0), R\nabla_{e_k}\xi(0)\rangle\rangle_\rho \\ &= \langle\langle T^*R\nabla_{e_j}\xi(0), R\nabla_{e_k}\xi(0)\rangle\rangle_\rho \\ &= \langle\langle \nabla_{e_j}\xi(0), T\nabla_{e_k}\xi(0)\rangle\rangle_\rho \\ &= Q_{kj}. \end{aligned}$$

So Q and thus the diffusion coefficient D are symmetric matrices. $\square$

Acknowledgments: This research has been supported by a Marie Curie Fellowship of the European Community Programme “Improving Human Potential” under the contract number HPMF-CT-2002-01610.

References

- [1] Chung, C.C., Landim, C., Olla, S. : *Equilibrium Fluctuations of Asymmetric Simple Exclusion Processes in Dimension $d \geq 3$* . Probab. Th. Rel. Fields **119** (2001), 381–409.
- [2] Esposito, R., Marra, R., Yau H.T. : *Diffusive Limit of Asymmetric Simple Exclusion*. Rev. Math. Phys. **6** (1994), 1233–1267.
- [3] Landim, C., Olla, S., Yau, H.T. : *First Order Correction for the Hydrodynamic Limit of Asymmetric Simple Exclusion Processes in Dimension $d \geq 3$* . Comm. Pure Appl. Math. **50** (1997), 149–203.
- [4] Landim, C., Olla, S., Yau, H.T. : *Some Properties of the Diffusion Coefficient for Asymmetric Simple Exclusion Processes*. Ann. Probab. **24** (1996), 1779–1807.
- [5] Landim, C., Olla, S., Varadhan, S.R.S. : *On Viscosity and Fluctuation-Dissipation in Exclusion Processes.*, J. Stat. Phys. **115** (2004), no1-2, 323–363.
- [6] Landim, C., Sued, M., Valle, G. : *Hydrodynamic Limit of Asymmetric Exclusion Process Under Diffusive Scaling in $d \geq 3$* , Commun. Math. Phys. **249** (2004), 215–247.
- [7] Xu, L. : *Diffusive Limit for a Lattice Gas with Short Range Interactions*. Ph.D. Thesis, New York University.